

## **Denaturation of protein by chlorine dioxide: oxidative modification of tryptophan and tyrosine residues**

Ogata N.

*Biochemistry* 46, 4898-4911 (2007).

<和文タイトル>

二酸化塩素によるタンパク質の変性：トリプトファンとチロシン残基の酸化的修飾

[Abstract]

Oxychlorine compounds, such as hypochlorous acid (HOCl) and chlorine dioxide (ClO<sub>2</sub>), have potent antimicrobial activity. Although the biochemical mechanism of the antimicrobial activity of HOCl has been extensively investigated, little is known about that of ClO<sub>2</sub>. Using bovine serum albumin and glucose-6-phosphate dehydrogenase of *Saccharomyces cerevisiae* as model proteins, here I demonstrate that the antimicrobial activity of ClO<sub>2</sub> is attributable primarily to its protein-denaturing activity. By solubility analysis, circular dichroism spectroscopy, differential scanning calorimetry, and measurement of enzymatic activity, I demonstrate that protein is rapidly denatured by ClO<sub>2</sub> with a concomitant decrease in the concentration of ClO<sub>2</sub> in the reaction mixture. Circular dichroism spectra of the ClO<sub>2</sub>-treated proteins show a change in ellipticity at 220 nm, indicating a decrease in  $\alpha$ -helical content. Differential scanning calorimetry shows that transition temperature and endothermic transition enthalpy of heat-induced unfolding decrease in the ClO<sub>2</sub>-treated protein. The enzymatic activity of glucose-6-phosphate dehydrogenase decreases to 10% within 15 s of treatment with 10  $\mu$ M ClO<sub>2</sub>. Elemental analyses show that oxygen, but not chlorine, atoms are incorporated in the ClO<sub>2</sub>-treated protein, providing direct evidence that protein is oxidized by ClO<sub>2</sub>. Furthermore, mass spectrometry and nuclear magnetic resonance spectroscopy show that tryptophan residues become *N*-formylkynurenine and tyrosine residues become 3,4-dihydroxyphenylalanine (DOPA) or 2,4,5-trihydroxyphenylalanine (TOPA) in the ClO<sub>2</sub>-treated proteins. Taking these results together, I conclude that microbes are inactivated by ClO<sub>2</sub> owing to denaturation of constituent proteins critical to their integrity and/or function, and that this denaturation is caused primarily by covalent oxidative modification of their tryptophan and tyrosine residues.

Reproduced with permission from *Biochemistry* 2007, 46, 4898-4911. Copyright 2007  
American Chemical Society.