Read this package insert carefully before using this product. Keep this package insert properly so that you can refer to it in case of necessity.



# **Supplement Facts**

Serving Size 3 Pills

	Amount Per Serving	% Daily Value
Phellodendron Bark	100mg	†
Citrus Unshiu Peel	100mg	†
Gambir Leaves	67mg	†
Licorice Root	50mg	†
T Daily Value not Established		

I Dally value not Established.

Other Ingredients :

Water, Beechwood Creosote, Glycerin, Calcium Carboxymethylcellulose, Cinnamon Bark

Uses:

SEIROGAN Pills is a herbal dietary supplement to help regulate digestive system to maintain normal bowel functions.\*

#### Directions:

Take 3 pills after a meal with water up to 3 times a day as needed. For children 5-10 years of age, the recommended serving is 1 pill. For age 11-14 years, the recommended serving is 2 pills. Not recommended for children under 5 years of age.

#### Warnings:

#### 請仔細閱讀使用説明書并保管好,以備需要時參考。



<b>成 份</b> 1次服用量 3 粒		
	1次服用量	1日攝取量%
黄柏粉	100mg	+
陳皮粉	100mg	†
阿仙藥粉	67mg	†
甘草粉	50mg	†
<b>†</b> 1日攝取量未定。		

#### 其他成分:

常水,木餾油,濃甘油,羧甲基纖維素鈣,桂皮粉

## 用途:

正露丸爲藥草營養輔助劑,能幫助維持消化系統及正常的腸道功能。\*

#### 用法·用量:

成人每次3粒,每日按需要可服用至3次,飯後以水服之。5至10歲兒童建議 每次服用1粒。11至14歲兒童建議每次服用2粒。建議未滿5歲兒童不要服用。

## 注意事項:

1. 孕產婦或授乳期的婦女, 請與醫護人員咨商後服用。

- 1. Pregnant or lactating women should not take this product without consulting their healthcare professionals. 2. Don't use this product to persons with a history of allergic symptoms to medicines or
- cosmetics.
- 3. Do not use other products claiming antidiarrheal effect while using this product.
- 4. Do not use this product continuously over a long-term period.
- 5. Do not exceed recommended dosage.
- 6. The following persons should consult with healthcare professionals before using this product: persons who are under treatment by a physician or a healthcare professional.
- persons with severe acute diarrhea; diarrhea accompanying fever; diarrhea accompanying bloody stools, mucous stools or symptoms of abdominal pain, abdominal distension or nausea; diabetics; hypertension or hypokalemia.
- persons who or whose family members have allergic disposition.
- persons with hepatic or renal disorders.
- elderly persons.
- persons taking thiazide diuretics, stimulant laxatives, cardiac glycocides or cortisol.
- 7. Discontinue to take this product in case the following symptoms occur after its use: Rash/redness, itch, edema, nausea/vomiting, constipation, anorexia, gastric discomfort, dizziness, headache, systemic dullness, jaundice due to hepatic dysfunction, etc.
- 8. Discontinue to use this product if symptoms are not improved even after several days of its use. 9. Keep out of reach of children.
- 10. For your protection, do not use if cap seal is broken or missing.
- 11. Do not transfer this product to another container in order to avoid misuse and maintain the quality.
- 12. Do not mix this product with other products.
- 13. If this product accidentally gets into eyes, wash immediately with water and visit an ophthalmologist.
- 14. If this product accidentally attaches to the skin, wash thoroughly with soap and hot water.

*Other Information :* 

Store in a cool place away from direct sunlight.

\* This statement has not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

#### *Manufactured by:*

## *Distributed by:*

TAIKO PHARMACEUTICAL CO., LTD. 3-34-14 Uchihonmachi, Suita, Osaka, Japan

PRINCE of PEACE ENTERPRISES, INC.® 3536 Arden Road Hayward, CA 94545-3908, U. S. A. Toll free # : 1 (800) 732-2328

- 2. 請不要給過去對藥品或化妝品有過敏癥狀之患者服用。
- 3. 請不要同時服用其他止瀉藥品。
- 4. 請不要長期持續服用。
- 5. 請按説明書中指示的劑量服用。
- 6. 服用前有以下癥狀者請向醫護人員請示。
  - 正在接受醫師的治療者。
  - ■具有劇烈急性腹瀉;伴有發燒癥狀的腹瀉;伴有血便,水便或腹痛,膨 脹感或惡心感的腹瀉;糖尿病;高血壓或低鉀血癥者。
  - 個人或家族具有過敏性者。
  - 肝, 腎功能不全者。
  - ■高齡者。
  - ■正在服用噻嗪類利尿藥,興奮性緩瀉藥,强心甙或皮質醇者。
- 7. 服用後若出現以下癥狀請停止服用:
  - 出疹/發紅,發癢,水腫,惡心/嘔吐,便秘,厭食,胃失常,頭暈,頭痛, 系統性的遲鈍, 肝機能不良所致黄疸等
- 8. 如服用數天而癥狀持續,請停止服用。
- 9. 請保管於小兒不易觸及之處。
- 10. 瓶蓋封印有破損或無包裝時, 請不要服用。
- 11. 爲避免誤用與保持品質,請勿將本品换置於其他容器中。
- 12. 請勿將其他的藥物混入本品。
- 13. 本品誤入眼睛時立即以水沖洗并求診於眼科醫生。
- 14. 本品誤與皮膚附着時可用肥皂與温水沖洗。

## 其他情報:

請保存于冷暗處,避免陽光照射。

\*此叙述未經美國食品藥物管理局之評估。本品非用於診斷,治療,醫治 或預防疾病所用。

#### 生産者: 🚺 大幸薬品株式会社 日本國大阪府吹田市内本町3丁目34番14號 經銷商:美國大子行 ↔

3536 Arden Road Hayward, CA 94545-3908, U. S. A. 消費熱線: 1 (800) 732-2328